

CLAIMS

1. A peritoneal dialysate containing a taurine compound along with an electrolyte and an alkalizer.

5 2. The peritoneal dialysate according to claim 1, containing the taurine compound in an amount of 0.01 to 5 w/v%.

3. The peritoneal dialysate according to claim 1, containing the taurine compound in an amount of 0.01 to 3.5 w/v%.

10 4. The peritoneal dialysate according to claim 1, containing the taurine compound in an amount of 0.01 to 1.5 w/v%.

5. The peritoneal dialysate according to claim 1, wherein the alkalizer is a lactate, a citrate, or a bicarbonate, and the electrolyte is sodium ion, calcium ion, magnesium ion, or chloride ion.

15 6. The peritoneal dialysate according to claims 1 or 5, having an osmotic pressure of 280 to 680 mOsm.

7. The peritoneal dialysate according to claims 1 or 5, having an osmotic pressure of 280 to 510 mOsm.

20 8. The peritoneal dialysate according to any one of claims 1 to 7, wherein the pH upon use is adjusted to a value of 6.0 to 7.5.

9. The peritoneal dialysate according to any one of claims 1 to 8, provided in a one-compartment container.

25 10. A peritoneal dialysate, containing 0.01 to 5 w/v% of taurine, 25 to 45 mEq/L of sodium lactate, 80 to 150 mEq/L of sodium ion, 0 to 3 mEq/L of potassium ion, 0.5 to 5 mEq/L of calcium ion, 0.1 to 2.0 mEq/L of magnesium ion, 80 to 110 mEq/L of chloride ion, and 0 to 4 w/v% of glucose and having a pH of 6.0 to 7.5 upon use.

30 11. The peritoneal dialysate according to claim 10, containing taurine in an amount of 0.01 to 3.5 w/v%.

12. The peritoneal dialysate according to claim 10, containing a taurine compound in an amount of 0.01 to 1.5 w/v%.

13. A peritoneal dialysate, containing 0.01 to 5 w/v% of

taurine, 25 to 45 mEq/L of sodium lactate, 80 to 150 mEq/L of sodium ion, 0 to 3 mEq/L of potassium ion, 0.5 to 5 mEq/L of calcium ion, 0.1 to 2.0 mEq/L of magnesium ion, and 80 to 110 mEq/L of chloride ion and having a pH of 6.0 to 7.5.

5 14. The peritoneal dialysate according to claim 13, containing taurine in an amount of 0.1 to 3.5 w/v%.

 15. The peritoneal dialysate according to claim 13, containing a taurine compound in an amount of 0.5 to 1.5 w/v%.